



THREATENED SPECIES HAZARD REDUCTION LIST – PART 3 – THREATENED ECOLOGICAL COMMUNITIES

Endangered ecological community	Listed on TSC Act	Listed on EPBC Act	Species Specific Conditions relating to the use of Fire	Species Specific Conditions relating to Mechanical Forms of Hazard Reduction
<i>Acacia loderi</i> shrublands	Endangered Ecological Community		No fire more than once every 6 years	No slashing, trittering or tree removal
<i>Acacia melvillei</i> Shrubland in the Riverina and Murray-Darling Depression bioregions	Endangered Ecological Community		No fire more than once every 20 years	No slashing, trittering or tree removal
Agnes Banks Woodland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
<i>Allocasuarina luehmannii</i> Woodland in the Riverina and Murray-Darling Depression Bioregions	Endangered Ecological Community	Endangered	No fire	No slashing, trittering or tree removal
Araluen Scarp Grassy Forest in the South East Corner Bioregion	Endangered Ecological Community		No fire more than once every 10 years	Slashing, but no trittering or tree removal
Artesian Springs Ecological Community	Endangered Ecological Community	Endangered	No fire	No slashing, trittering or tree removal
Bangalay Sand Forest of the Sydney Basin and South East Corner bioregions	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Ben Halls Gap National Park Sphagnum Moss Cool Temperate Rainforest	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Blue Gum High Forest in the Sydney Basin Bioregion	Critically Endangered Ecological Community	Critically Endangered	Low intensity fire only. No fire more than once every 15 years	No slashing, trittering or tree removal
Blue Mountains Shale Cap Forest in the Sydney Basin Bioregion	Endangered Ecological Community	Critically Endangered	No fire more than once every 10 years	No slashing, trittering or tree removal
Blue Mountains Swamps in the Sydney Basin Bioregion	Vulnerable Ecological Community	Endangered	No fire more than once every 12 years	No slashing, trittering or tree removal
Brigalow within the Brigalow Belt South, Nandewar and Darling Riverine Plains Bioregions	Endangered Ecological Community	Endangered	No fire more than once every 6 years	No slashing, trittering or tree removal
Brigalow-Gidgee woodland/shrubland in the Mulga Lands and Darling Riverine Plains Bioregions	Endangered Ecological Community	Endangered	No fire more than once every 15 years	No slashing, trittering or tree removal
Brogo Wet Vine Forest in the South East Corner Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Byron Bay Dwarf Graminoid Clay Heath Community	Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
<i>Cadellia pentastylis</i> (Ooline) community in the Nandewar and Brigalow Belt South Bioregions	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Carbeen Open Forest Community in the Darling Riverine Plains and Brigalow Belt South Bioregions	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Castlereagh Scribbly Gum Woodland in the Sydney Basin Bioregion	Vulnerable Ecological Community	Endangered	No fire more than once every 12 years	No slashing, trittering or tree removal
Castlereagh Swamp Woodland Community	Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
Central Hunter Grey Box-Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions	Endangered Ecological Community		No fire more than once every 8 years	No slashing, trittering or tree removal
Central Hunter Ironbark-Spotted Gum-Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions	Endangered Ecological Community		No fire more than once every 8 years	No slashing, trittering or tree removal
Coastal Cypress Pine Forest in the New South Wales North Coast Bioregion	Endangered Ecological Community		No fire more than once every 20 years	No slashing, trittering or tree removal
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin	Endangered Ecological Community	Vulnerable	No fire	No slashing, trittering or tree removal

Please Note: This List is to be applied with reference to the "Rules and Notes for the implementation of the Threatened Species Hazard Reduction List for the Bush Fire Environmental Assessment Code". This List was compiled on 25th September 2013.

Endangered ecological community and South East Corner Bioregions	Listed on TSC Act	Listed on EPBC Act	Species Specific Conditions relating to the use of Fire	Species Specific Conditions relating to Mechanical Forms of Hazard Reduction
Cooks River/Castlereagh Ironbark Forest in the Sydney Basin Bioregion	Endangered Ecological Community	Critically Endangered	No fire more than once every 7 years	No slashing, trittering or tree removal
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Penepplain and Mulga Lands Bioregion	Endangered Ecological Community	Endangered	No fire	No slashing, trittering or tree removal
Cumberland Plain Woodland in the Sydney Basin Bioregion	Critically Endangered Ecological Community	Critically Endangered	No fire more than once every 7 years	No slashing, trittering or tree removal
Dry Rainforest of the South East Forests in the South East Corner Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Duffys Forest Ecological Community in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Eastern Suburbs Banksia Scrub in the Sydney Basin Bioregion	Endangered Ecological Community	Endangered	No fire more than once every 10 years	No slashing, trittering or tree removal
Elderslie Banksia Scrub Forest in the Sydney Basin Bioregion	Critically Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Endangered Ecological Community		No fire more than once every 15 years	No slashing, trittering or tree removal
Genowlan Point <i>Allocasuarina nana</i> Heathland	Endangered Ecological Community		No fire more than once every 20 years	No slashing, trittering or tree removal
Grey Box-Grey Gum Wet Sclerophyll Forest in the NSW North Coast Bioregion	Endangered Ecological Community		No fire more than once every 30 years	No slashing, trittering or tree removal
<i>Halosarcia lylei</i> low open-shrubland in the Murray Darling Depression Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Howell Shrublands in the New England Tableland and Nandewar Bioregions	Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
Hunter Floodplain Red Gum Woodland in the NSW North Coast and Sydney Basin Bioregions	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions	Endangered Ecological Community		No fire more than once every 25 years	Slashing, but no trittering or tree removal
Hunter Valley Foothills Slaty Gum Woodland in the Sydney Basin Bioregion	Vulnerable Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Hunter Valley Vine Thicket in the NSW North Coast and Sydney Basin Bioregions	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Hunter Valley Weeping Myall Woodland of the Sydney Basin Bioregion	Endangered Ecological Community	Critically Endangered	No fire	No slashing, trittering or tree removal
Hygrocybeae Community of Lane Cove Bushland Park in the Sydney Basin Bioregion	Critically Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Illawarra Lowlands Grassy Woodland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 5 years for grassy woodland sub-community, and no more than once every 25 years for the moist forest sub-community	No slashing, trittering or tree removal
Illawarra Subtropical Rainforest in the Sydney Basin Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Penepplain, Nandewar and Brigalow Belt South Bioregions	Endangered Ecological Community	Endangered	No fire more than once every 15 years	No slashing, trittering or tree removal
Kincumber Scribbly Gum Forest in the Sydney Basin Bioregion	Critically Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
Kurnell Dune Forest in the Sutherland Shire and City of Rockdale	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Kurri Sand Swamp Woodland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal

Please Note: This List is to be applied with reference to the "Rules and Notes for the implementation of the Threatened Species Hazard Reduction List for the Bush Fire Environmental Assessment Code". This List was compiled on 25th September 2013.

Endangered ecological community	Listed on TSC Act	Listed on EPBC Act	Species Specific Conditions relating to the use of Fire	Species Specific Conditions relating to Mechanical Forms of Hazard Reduction
Littoral Rainforest in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Endangered Ecological Community	Critically Endangered	No fire	No slashing, trittering or tree removal
Low woodland with heathland on indurated sand at Norah Head	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Lower Hunter Spotted Gum-Ironbark Forest in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 8 years	No slashing, trittering or tree removal
Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions	Vulnerable Ecological Community		No fire	No slashing, trittering or tree removal
Lowland Grassy Woodland in the South East Corner Bioregion	Endangered Ecological Community	Critically Endangered	No fire more than once every 8 years	No slashing, trittering or tree removal
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Endangered Ecological Community	Critically Endangered	No fire	No slashing, trittering or tree removal
Lowland Rainforest on Floodplain in the New South Wales North Coast Bioregion	Endangered Ecological Community	Critically Endangered	No fire	No slashing, trittering or tree removal
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking <i>Triodia</i> , in the NSW South Western Slopes Bioregion	Critically Endangered Ecological Community		No fire more than once every 25 years	No slashing, trittering or tree removal
Maroota Sands Swamp Forest	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Marsh Club-rush sedgeland in the Darling Riverine Plains Bioregion	Critically Endangered Ecological Community		No fire	No slashing, trittering or tree removal
McKies Stringybark/Blackbutt Open Forest in the Nandewar and New England Tableland Bioregions	Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
<i>Melaleuca armillaris</i> Tall Shrubland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Milton Ulladulla Subtropical Rainforest in the Sydney Basin Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Moist Shale Woodland in the Sydney Basin Bioregion	Endangered Ecological Community	Critically Endangered	No fire more than once every 10 years	No slashing, trittering or tree removal
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps bioregions	Endangered Ecological Community	Endangered	No fire	No slashing, trittering or tree removal
Mount Gibraltar Forest in the Sydney Basin Bioregion	Endangered Ecological Community	Endangered	No fire more than once every 7 years	No slashing, trittering or tree removal
Mt Canobolas <i>Xanthoparmelia</i> Lichen Community	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes bioregions	Endangered Ecological Community	Endangered	No fire more than once every 25 years	No slashing, trittering or tree removal
Native Vegetation on Cracking Clay Soils of the Liverpool Plains	Endangered Ecological Community	Critically Endangered	No fire more than once every 2 years	
New England Peppermint (<i>Eucalyptus nova-anglica</i>) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Critically Endangered Ecological Community	Critically Endangered	No fire more than once every 10 years	No slashing, trittering or tree removal
Newnes Plateau Shrub Swamp in the Sydney Basin Bioregion	Endangered Ecological Community	Endangered	No fire more than once every 11 years	No slashing, trittering or tree removal
O'Hares Creek Shale Forest	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Pittwater and Wagstaffe Spotted Gum Forest in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Porcupine Grass - Red Mallee - Gum Coolabah hummock grassland/low sparse woodland in the Broken Hill Complex Bioregion	Critically Endangered Ecological Community		No fire more than once every 25 years	No slashing, trittering or tree removal
Quorrobolong Scribbly Gum Woodland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Ribbon Gum-Mountain Gum-Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Endangered Ecological Community		No fire more than once every 15 years	No slashing, trittering or tree removal
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Endangered Ecological Community		No fire more than once every 11 years	No slashing, trittering or tree removal

Please Note: This List is to be applied with reference to the "Rules and Notes for the implementation of the Threatened Species Hazard Reduction List for the Bush Fire Environmental Assessment Code". This List was compiled on 25th September 2013.

Endangered ecological community	Listed on TSC Act	Listed on EPBC Act	Species Specific Conditions relating to the use of Fire	Species Specific Conditions relating to Mechanical Forms of Hazard Reduction
Robertson Basalt Tall Open-forest in the Sydney Basin Bioregion	Endangered Ecological Community	Endangered	No fire more than once every 25 years, and of low intensity	No slashing, trittering or tree removal
Robertson Rainforest in the Sydney Basin Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Sandhill Pine Woodland in the Riverina, Murray-Darling Depression and NSW South Western Slopes bioregions	Endangered Ecological Community		No fire more than once every 15 years	No slashing, trittering or tree removal
Semi-evergreen Vine Thicket in the Brigalow Belt South and Nandewar Bioregions	Endangered Ecological Community	Endangered	No fire	No slashing, trittering or tree removal
Shale Gravel Transition Forest in the Sydney Basin Bioregion	Endangered Ecological Community	Critically Endangered	No fire more than once every 7 years	No slashing, trittering or tree removal
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Critically Endangered Ecological Community	Critically Endangered	No fire more than once every 7 years	No slashing, trittering or tree removal
Southern Highlands Shale Woodlands in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Southern Sydney sheltered forest on transitional sandstone soils in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Endangered Ecological Community		No fire more than once every 15 years	No slashing, trittering or tree removal
Sun Valley Cabbage Gum Forest in the Sydney Basin Bioregion	Critically Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Sydney Freshwater Wetlands in the Sydney Basin Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
Sydney Turpentine-Ironbark Forest	Endangered Ecological Community	Critically Endangered	No fire more than once every 10 years	No slashing, trittering or tree removal
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Endangered Ecological Community		No fire more than once every 15 years	No slashing, trittering or tree removal
The Shorebird Community occurring on the relict tidal delta sands at Taren Point	Endangered Ecological Community		No fire	No slashing, trittering or tree removal
<i>Themeda</i> grassland on seacliffs and coastal headlands in the NSW North Coast, Sydney Basin and South East Corner Bioregions	Endangered Ecological Community		No fire more than once every 7 years	No slashing, trittering or tree removal
Umina Coastal Sandplain Woodland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Upland Wetlands of the Drainage Divide of the New England Tableland Bioregion	Endangered Ecological Community	Endangered	No fire more than once every 6 years	No slashing, trittering or tree removal
Warkworth Sands Woodland in the Sydney Basin Bioregion	Endangered Ecological Community		No fire more than once every 10 years	No slashing, trittering or tree removal
Western Sydney Dry Rainforest in the Sydney Basin Bioregion	Endangered Ecological Community	Critically Endangered	No fire	No slashing, trittering or tree removal
White Box Yellow Box Blakely's Red Gum Woodland	Endangered Ecological Community	Critically Endangered	No fire more than once every 5 years	Slashing, but no trittering or tree removal
White Gum Moist Forest in the NSW North Coast Bioregion	Endangered Ecological Community		No fire	No slashing, trittering or tree removal

Please Note: This List is to be applied with reference to the "Rules and Notes for the implementation of the Threatened Species Hazard Reduction List for the Bush Fire Environmental Assessment Code". This List was compiled on 25th September 2013.